ST AVAILABLE COPY







PuhMed	Nucketida	Pipled	Genome	Structure	fic	pSot	Yanonoary	OMIM
Search PubM	1ed ▼ fo	r Limits	Preview/Inc	Hav	History	Go	Cliphoard	Deta
About Entrez	Dis	ļ			Save	Text	Clip Add	Order
Entrez PubMed Overviow						Rela	ated Articles, .NEF	Books.

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Quories
LinkOut

New/Noteworthy

Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts Clinical Trials.gov PubMed Central

Cubby

Privacy Policy

Predicting the topology of eukaryotic membrane proteins.

Sipos L, von Heijne G.

Department of Theoretical Physics, Royal Institute of Technology, Stockholm, Sweden.

We show that the so-called 'positive inside' rule, i.e. the observation that positively charged amino acids tend to be more prevalent in cytoplasmic than in extra-cytoplasmic segments in transmembrane proteins [von Heijne, G. (1986) EMBO J. 5, 3021-3027], seems to hold for all polar segments in multi-spanning eukaryotic membrane proteins irrespective of their position in the sequence and hence can be used in conjunction with hydrophobicity analysis to predict their transmembrane topology. Further, as suggested by others, we confirm that the net charge difference across the first transmembrane segment correlates well with its orientation [Hartmann, E., Rapoport, T. A. and Lodish H. F. (1989) Proc. Natl Acad. Sci. USA 86, 5786-5790], and that the overall amino-acid composition of long polar segments can also be used to predict their cytoplasmic or extracytoplasmic location [Nakashima, H. and Nishikawa, K. (1992) FEBS Lett. 303, 141-146]. We present an approach to the topology prediction problem for eukaryotic membrane proteins based on a combination of these methods.

PMID: 8099327 [PubMed - indexed for MEDLINE]

Display	Abstract	Sort	<u>-</u>	Save	Text	Clip Add	Order

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer